

# Psychiatric Comorbidity and Psychosocial Factors Matter in Heart Failure

Ching-Yen Chen<sup>1,2</sup> and Chun-Li Wang<sup>2,3</sup>

Comment on “Heart failure functional class associated with depression severity but not anxiety severity” by Celik E. et al. *Acta Cardiologica Sinica* 2016;32:55-61.

The incidence and prevalence of heart failure (HF) continues to increase despite the advances in therapies that have occurred.<sup>1</sup> The impact of psychiatric co-morbidity and psychosocial factors on HF has been studied extensively. Many studies have focused on the link between depression and HF.<sup>2-5</sup> While depression in patients with HF is the psychiatric problem most commonly discussed,<sup>6</sup> anxiety is often present concurrently with depression, and may be associated with poor prognosis in patients with HF. However, only a few studies have examined the association between anxiety and HF, and the results of these studies have been inconsistent.<sup>7,8</sup> Particularly, 8-16% of patients with HF suffer from distressful anxiety, such as generalized anxiety disorder or panic disorder.<sup>9,10</sup> Overall anxiety levels are 60% higher in HF patients than in the healthy elderly population.<sup>11</sup>

Given that anxiety disorders differ in types and symptoms, assessing the severity of anxiety may be more difficult than assessing the severity of depression. Considering its brevity, simplicity, and presumed ability to measure general anxiety, the Beck Anxiety Inventory (BAI) might be a good candidate for clinical use as a severity indicator.<sup>12</sup> However, the value of BAI has been

disputed for its focus on psycho-physiological symptoms linked to panic.<sup>13,14</sup> Arguably, the instrument seems to reflect the severity of depression, but is not a suitable scale to effectively differentiate between anxiety and depression.<sup>12</sup>

Importantly, many studies have shown that depression, but not anxiety, was associated with poor outcomes in HF patients who were hospitalized<sup>3,15</sup> or had a low ejection fraction (< 35%).<sup>9,16</sup> Miyuki et al. observed that anxiety was independently associated with adverse outcomes in patients with mild HF.<sup>17</sup> Even anxiety has been linked with prognosis, morbidity, and functional status in patients with HF, although only a few studies have investigated the determinants of anxiety in patients with HF. The distressed (type-D) personality may be another important, but easily ignored, determinant of anxiety in HF patients.<sup>18</sup> Type-D personality is referred to as the combined tendency to experience negative emotions (high negative affectivity) and to inhibit self-expression in social interactions (high social inhibition).<sup>19</sup> Additionally, type-D personality has been associated with worse prognosis and impaired quality of life in patients with coronary artery disease.<sup>20</sup> Furthermore, type-D personality is independently linked with an increased risk of depression and anxiety in patients with HF.<sup>21</sup>

In the study of Miyuki et al., depression was associated with living alone and poor social support rather than medical factors among patients with HF.<sup>17</sup> Social factors were important predictors for depression and anxiety, suggesting that the establishment of a social support system should be an essential element in HF treatment. Patients with chronic HF often have cognitive impairment, particularly in memory, psychomotor speed,

Received: May 8, 2015 Accepted: July 23, 2015

<sup>1</sup>Department of Psychiatry, Chang Gung Memorial Hospital, Linkou; <sup>2</sup>School of Medicine, Chang Gung University, Taoyuan; <sup>3</sup>Division of Cardiology, Department of Internal Medicine, Chang Gung Memorial Hospital, Linkou, Taoyuan, Taiwan.

Address correspondence and reprint requests to: Dr. Chun-Li Wang, Division of Cardiology, Department of Internal Medicine, Chang Gung Memorial Hospital, Linkou, Taoyuan, No. 5, Fushin St. Kwei-Shan District, Taoyuan, Taiwan. Tel: 886-3-328-1200 ext. 8162; E-mail: wang3015@cgmh.org.tw

and executive function.<sup>22</sup> Cognitive impairment in HF patients is related to grey matter loss which has been revealed by a brain imaging study.<sup>23</sup> Cognitive impairment is multi-dimensional, and poor concentration is one of its several possible presentations. Poor concentration is also included in depression and anxiety symptoms cluster. Therefore, cognitive deficit in HF patients is associated with both neurological and psychiatric factors.

In the current study, depression severity, but not anxiety severity, is positively associated with HF symptom class.<sup>24</sup> However, several points do need to be addressed. First, this cross-sectional study can only show the severity of depression increases with HF symptom class, but cannot show that depression increases the risk of HF. Second, the definition and selected psychometric scale for determining the severity of anxiety may affect the relationship between anxiety and HF symptom class. The self-report questionnaires such as the BAI might not be sufficiently precise to make a psychiatric diagnosis of anxiety, and differentiate between anxiety and depression in patients with HF. Third, symptoms of HF, treatment and timing of evaluation may influence the values of BAI and the observed relationship.

Highly prevalent psychiatric co-morbidity and psycho-social factors are commonly observed in patients with HF. This field deserves additional attention in order to develop a comprehensive evidence-based assessment tool that comprises depression, anxiety, personality trait, cognitive function, and psycho-social stresses for patients with HF. With this tool, clinicians may better identify subjects at increased risk, and adjust therapeutic strategies accordingly to improve patient outcomes.

## REFERENCES

1. Chapa DW, Akintade B, Son H, et al. Pathophysiological relationships between heart failure and depression and anxiety. *Crit Care Nurse* 2014;34:14-23.
2. Faris R, Purcell H, Henein MY, Coats AJ. Clinical depression is common and significantly associated with reduced survival in patients with non-ischemic heart failure. *Eur J Heart Fail* 2002; 4:541-51.
3. Jiang W, Kuchibhatla M, Cuffe MS, et al. Prognostic value of anxiety and depression in patients with chronic heart failure. *Circulation* 2004;110:3452-6.
4. Sherwood A, Blumenthal JA, Trivedi R, et al. Relationship of depression to death or hospitalization in patients with heart failure. *Arch Intern Med* 2007;167:367-73.
5. Skotzko CE, Krichten C, Zietowski G, et al. Depression is common and precludes accurate assessment of functional status in elderly patients with congestive heart failure. *J Card Fail* 2000; 6:300-5.
6. Yohannes AM, Willgoss TG, Baldwin RC, Connolly MJ. Depression and anxiety in chronic heart failure and chronic obstructive pulmonary disease: prevalence, relevance, clinical implications and management principles. *Int J Geriatr Psychiatry* 2010;25:1209-21.
7. MacMahon KM, Lip GY. Psychological factors in heart failure: a review of the literature. *Arch Intern Med* 2002;162:509-16.
8. Konstam V, Moser DK, De Jong MJ. Depression and anxiety in heart failure. *J Card Fail* 2005;11:455-63.
9. Haworth JE, Moniz-Cook E, Clark AL, et al. Prevalence and predictors of anxiety and depression in a sample of chronic heart failure patients with left ventricular systolic dysfunction. *Eur J Heart Fail* 2005;7:803-8.
10. Griez EJ, Mammars N, Loirat JC, et al. Panic disorder and idiopathic cardiomyopathy. *J Psychosom Res* 2000;48:585-7.
11. Denollet J, Brutsaert DL. Personality, disease severity, and the risk of long-term cardiac events in patients with a decreased ejection fraction after myocardial infarction. *Circulation* 1998; 97:167-73.
12. Beck AT, Epstein N, Brown G, Steer RA. An inventory for measuring clinical anxiety: psychometric properties. *J Consult Clin Psychology* 1988;56:893-7.
13. Cox BJ, Cohen E, Dorenfeld DM, Swinson RP. Does the Beck Anxiety Inventory measure anything beyond panic attack symptoms? *Behav Res Ther* 1996;34:949-54.
14. Leyfer OT, Ruberg JL, Woodruff-Borden J. Examination of the utility of the Beck Anxiety Inventory and its factors as a screener for anxiety disorders. *J Anxiety Disord* 2006;20:444-58.
15. Vaccarino V, Kasl SV, Abramson J, Krumholz HM. Depressive symptoms and risk of functional decline and death in patients with heart failure. *J Am Coll Cardiol* 2001;38:199-205.
16. Junger J, Schellberg D, Muller-Tasch T, et al. Depression increasingly predicts mortality in the course of congestive heart failure. *Eur J Heart Fail* 2005;7:261-7.
17. Miyuki TM, Norihiro K, Akiko C, et al. Anxiety and poor social support are independently associated with adverse outcomes in patients with mild heart failure. *Circ J* 2009;73:280-7.
18. Schiffer AA, Pedersen SS, Broers H, et al. Type-D personality but not depression predicts severity of anxiety in heart failure patients at 1-year follow-up. *J Affect Disord* 2008;106:73-81.
19. Denollet J. DS14: standard assessment of negative affectivity, social inhibition, and type D personality. *Psychosom Med* 2005; 67:89-97.
20. Al-Ruzzeh S, Athanasiou T, Magoush O, et al. Predictors of poor mid-term health related quality of life after primary isolated coronary bypass grafting surgery. *Heart* 2005;91:1557-62.

21. van Gestel YR, Pedersen SS, van de Sande M, et al. Type-D personality and depressive symptoms predict anxiety 12 months post-percutaneous coronary interventions. *J Affect Disord* 2007; 103:197-203.
22. Sloan RS, Pressler SJ. Cognitive deficits in heart failure: recognition of vulnerability as a strange new world. *J Cardiovasc Nurs* 2009;24:241-8.
23. Almeida OP, Garrido GJ, Beer C, et al. Cognitive and brain changes associated with ischaemic heart disease and heart failure. *Eur Heart J* 2012;33:1769-76.
24. Celik E, Cay S, Sensoy B, et al. Heart failure functional class associated with depression severity but not anxiety severity. *Acta Cardiol Sin* 2016;32:55-61.

